

Compound-Specific Isotopic Fractionation Patterns Suggest Different Carbon Metabolisms among *Chloroflexus*-Like Bacteria in Hot-Spring Microbial Mats†

Marcel T. J. van der Meer,^{1*} Stefan Schouten,¹ Jaap S. Sinninghe Damsté,¹
Jan W. de Leeuw,¹ and David M. Ward²

Department of Marine Biogeochemistry and Toxicology, Royal Netherlands Institute for Sea Research, 1790 AB Den Burg, Texel, The Netherlands,¹ and Department of Land Resources and Environmental Sciences, Montana State University, Bozeman, Montana 59717²

Received 28 April 2003/Accepted 25 July 2003

Stable carbon isotope fractionations between dissolved inorganic carbon and lipid biomarkers suggest photoautotrophy by *Chloroflexus*-like organisms in sulfidic and nonsulfidic Yellowstone hot springs. Where co-occurring, cyanobacteria appear to cross-feed *Chloroflexus*-like organisms supporting photoheterotrophy as well, although the relatively small ¹³C fractionation associated with cyanobacterial sugar biosynthesis may sometimes obscure this process.

Chloroflexus aurantiacus and its phylogenetic relatives, which comprise a deeply branching kingdom-level lineage in the domain *Bacteria* (24), are major components of photosynthetic microbial mats in both sulfidic and nonsulfidic hot springs in Yellowstone National Park, Wyo. (3, 4, 5, 12, 23, 47). *C. aurantiacus*, the most studied representative of the green nonsulfur bacteria available in pure cultures, can grow heterotrophically by aerobic respiration, photoheterotrophically (using light to incorporate prerduced organic compounds), and photoautotrophically (using light to fix inorganic carbon) (26). Photoautotrophic metabolism by an obligately phototrophic relative of *C. aurantiacus*, the predominant phototroph in sulfidic hot-spring microbial mats, has been reported (12). However, based on culture (25, 26) and radiolabeling studies (2, 32), it has been hypothesized (44) that *Chloroflexus*-like organisms are mainly photoheterotrophic in mats where they live together with cyanobacteria. In such mats cyanobacteria are thought to be the main primary producers, which cross-feed reduced organic compounds to *Chloroflexus*-like organisms, thus supporting their photoheterotrophic metabolism. We exploited the distinctive lipid biomarkers of cyanobacteria and *Chloroflexus*-like organisms to test this hypothesis (39). If *Chloroflexus*-like organisms were purely photoheterotrophic in cyanobacterial mats, their biomarkers should have a $\delta^{13}\text{C}$ signature, similar to biomarkers of cyanobacteria. Lipids derived from cyanobacteria in hot-spring mats were found to have $\delta^{13}\text{C}$ values typical of the -20 to -25% fractionations relative to the inorganic carbon source expected of the Calvin cycle (27, 31). However, long-chain polyunsaturated alkenes (e.g., the $\text{C}_{31:3}$ alkene hentriacontatriene) and C_{30-37} wax esters that are typical of *Chloroflexus* (17, 35, 40) and mats containing *Chlo-*

roflexus relatives (9, 35, 37, 39) were approximately 10 to 15‰ enriched in ¹³C relative to cyanobacterial lipids (37, 39). Such values are expected based on autotrophic metabolism by *Chloroflexus*, which is known to use an inorganic carbon fixation pathway, the 3-hydroxypropionate pathway (15), that imparts an unusually heavy $\delta^{13}\text{C}$ signature to both biomass and lipids (16, 40). Thus, the enrichment in ¹³C of *Chloroflexus* lipids relative to cyanobacterial lipids in microbial mats pointed towards autotrophic growth of *Chloroflexus*-like organisms in hot-spring microbial mats (39).

Here, we examine an alternative explanation—*Chloroflexus*-like biomarkers could have heavy $\delta^{13}\text{C}$ signatures because they are cross-fed isotopically heavy fixed organic matter from cyanobacteria. Radiolabeling studies have shown that light-driven CO₂ fixation in hot-spring cyanobacterial mats leads mainly to production of polysaccharides (18, 22), presumably by cyanobacteria, and the polysaccharides are fermented during darkness to short-chain fatty acids known to be photoassimilated by *Chloroflexus*-like organisms (2, 23, 32). It has been shown that sugars may be significantly enriched in ¹³C relative to lipids in cyanobacteria and other Calvin cycle photoautotrophs (7, 43). Hence, we now have significantly expanded our previous study by investigating not only the $\delta^{13}\text{C}$ signatures of lipid biomarkers and biomass but also sugars and dissolved inorganic carbon (DIC) species to enable determination of isotopic fractionations. Furthermore, we included more representatives of mats where cyanobacteria and *Chloroflexus*-like organisms live together in different environments, thereby enabling us to better observe differences in isotopic fractionation patterns.

MATERIALS AND METHODS

Samples were taken from five different hot-spring microbial mats containing *Chloroflexus*-like bacteria with and without cyanobacteria located in Yellowstone National Park, Wyo. (Table 1). Samples for lipid analysis were frozen in the field and were kept frozen until lyophilization and lipid extraction. Samples for microscopy were stored on ice and directly analyzed after returning from the field. The presence of cyanobacteria and *Chloroflexus*-like bacteria was determined by phase contrast and autofluorescence microscopy. Water samples for sulfide anal-

* Corresponding author. Mailing address: Royal Netherlands Institute for Sea Research, Department of Marine Biogeochemistry and Toxicology, P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands. Phone: (31) 222-369565. Fax: (31) 222-319674. E-mail: mmeer@nioz.nl.

† NIOZ contribution no. 3630; journal series no. 2003-27, Montana Agricultural Experiment Station, Montana State University—Bozeman.

TABLE 1. General description of mat samples

Mat samples	Geothermal area	Temp (°C)	pH	[S ²⁻] (μM)	<i>Chloroflexus</i> -like	Cyanobacteria	Sampling date ^d
NMA source ^a	Mammoth Terraces	58.9	6.1	133	+	–	05-12-1995
NMA downstream ^a	Mammoth Terraces	56.3	6.4	19	+	+	05-12-1995
Tangerine Spring ^b	Mammoth Terraces	60	6.4	40	+	+	26-08-1999
Mushroom Spring ^c	Lower Geyser Basin	64	8.3	BD ^e	+	+	25-08-1997
Octopus Spring ^c	Lower Geyser Basin	58–64	8.3	BD	+	+	27-08-1997

^a Previously reported by van der Meer et al. (39).

^b On the shoulder of Orange Mound Spring along the Mammoth Upper Terrace loop road.

^c See Ward et al. (45) and Ramsing et al. (28) for location.

^d Day-month-year.

^e BD, below detection.

ysis were collected from above each mat, preserved in zinc acetate, and analyzed by the method of Cline (6). Inorganic carbon in water overflowing mats was trapped as BaCO₃ by increasing the pH of 200 ml of spring water to approximately pH 11 by adding a saturated NaOH solution (pH 13) and solid BaCl₂ (36). From the stable carbon isotopic composition of the BaCO₃, the isotopic composition of the CO₂ in the spring water was calculated using the temperature-dependent isotopic equilibrium equation of Mook et al. (21).

Lipids were extracted, derivatized, and analyzed by using gas chromatography, gas chromatography-mass spectrometry, and isotope-ratio-monitoring gas chromatography-mass spectrometry (33). Cell-associated sugars were analyzed by using approximately 20 to 160 mg of microbial mat residue after lipid extraction. This material was hydrolyzed, and sugar monomers were derivatized and analyzed by gas chromatography, gas chromatography-mass spectrometry, and isotope-ratio-monitoring gas chromatography-mass spectrometry (42). Stable carbon isotopic compositions of the bulk cell material and BaCO₃ were determined by automated on-line combustion (Carlo Erba CN analyser 1502 series) followed by conventional isotope ratio-mass spectrometry (Fisons optima [11]). The stable carbon isotope compositions are reported in the delta notation relative to the Vienna PeeDee Belemnite ¹³C standard.

RESULTS AND DISCUSSION

The mat samples studied are compared in Table 1. The New Mound Annex (NMA) mat, found in a sulfidic, high-carbonate hot spring at Mammoth Terraces, is comprised of *Chloroflexus* without cyanobacteria (39). The sulfide is thought to poison cyanobacteria but is utilized by *Chloroflexus* spp. in a photoautotrophic metabolism (12). All other mats are comprised of both cyanobacteria (*Synechococcus*) and *Chloroflexus*-like filamentous bacteria. The Tangerine Spring mat, like the downstream NMA mat (39), occurs in a slightly acidic, high-carbonate spring in the Mammoth Terraces group, where sulfide levels have been reduced due to its use by *Chloroflexus* spp. and other upstream sulfide-utilizing organisms. The Octopus Spring and Mushroom Spring mats occur in low-sulfide, alkaline silica-rich springs of the Lower Geyser Basin.

Lipid and sugar composition. The lipid compositions of the five mat samples are compared in Fig. 1 and Table 2, which also includes information for cultures of *C. aurantiacus* and its phylogenetic relative, *Roseiflexus castenholzii* (13), for reference. Lipids characteristic of *Chloroflexus*-like organisms were abundant in all mats. For instance, wax esters ranging from C₃₁ to C₃₇ were among the predominant lipids in all mats, with small-scale variation in distribution pattern among mat samples. Differences in the carbon skeletons of the wax esters between the environmental samples and the cultures may reflect physiological differences between cultivated and natural populations. For instance, the different mat systems do not contain monounsaturated wax esters as does *C. aurantiacus* but rather contain *iso*-branched wax esters (35, 39). Also, the mats

contain C₃₁ to C₃₇ wax esters, whereas *R. castenholzii* produces C₃₇ to C₄₀ wax esters (41). Long-chain (C₂₉₋₃₂) alkenes, predominantly hentriacontatriene (C_{31:3}) (38), were abundant in mats in Mammoth Terraces springs but were present at only trace levels in mats from the Lower Geyser Basin. This may reflect the predominance of *Chloroflexus* spp. in Mammoth Terraces (46) and more distantly related *Roseiflexus*-like organisms in springs of the Lower Geyser Basin (23, 30), since the latter organism lacks hentriacontatriene (41). C₁₇ *n*-alkane, a biomarker for cyanobacteria (34), was found in all mats containing cyanobacteria but was absent from the NMA source mat. More common lipids, such as C₁₅₋₁₈ fatty acids and C₁₇₋₁₈ alcohols, were also detected in all mats.

The sugar fractions of all mats contained arabinose, xylose, rhamnose, and glucose, the latter being dominant (approximately 30 to 80% of the sugar fraction). The sugar distribution was similar to that reported for *C. aurantiacus* (40).

Stable carbon isotopic compositions. The isotopic compositions of DIC, bulk biomass, lipids, and sugars for all mats are reported in Table 3. The values for DIC species determined for Tangerine Spring was within the range previously reported for other Mammoth Terraces hot springs (19). The values for Mushroom Spring were somewhat lower. Isotopic composition of bulk biomass ranged from –13 to –17‰, except for the NMA downstream and Tangerine Spring mats, which were isotopically lighter (–24 and –25‰, respectively). Isotopic compositions of specific lipids ranged from –8.9 to –36.3‰. Compounds known from or possibly contributed by cyanobacteria (e.g., C₁₇ *n*-alkane and C₁₆ and C₁₈ fatty acids, respectively) were isotopically lighter (–21.3 to –36.3‰) than biomarkers of *Chloroflexus*-like organisms (C_{31:3} and wax esters; –8.9 to –27.0‰). The isotopic composition of glucose ranged from –5.1 to –21.8‰. Isotopic compositions in cyanobacterial mats were generally more depleted in ¹³C in springs from Mammoth Terraces (Tangerine and NMA downstream) than in springs from the Lower Geyser Basin (Octopus Spring and Mushroom Spring).

Comparison of ¹³C fractionation in mats and cultures. Isotopic fractionations between bicarbonate, the principle DIC species, and bulk biomass, lipids, and glucose for all mat samples and *C. aurantiacus* are compared in Fig. 2. The ¹³C fractionations between biomass, lipids, and glucose relative to DIC are similar for the photoautotrophically grown *C. aurantiacus* culture and the *Chloroflexus* spp.-dominated NMA source pool microbial mat (Fig. 2). The smallest ¹³C fractionation relative to DIC (Δδ¹³C_{DIC}) was observed for glucose, followed by hen-




FIG. 1. Partial gas chromatograms of the total lipid extracts from the investigated microbial mats. 1, C_{17} *n*-alkane; 2, C_{15} fatty acid; 3, C_{16} fatty acid; 4, C_{17} fatty acid; 5, C_{17} alkanol; 6, $C_{18:1}$ fatty acid; 7, C_{18} fatty acid; 8, C_{18} alkanol; 9, $C_{31:3}$ alkene; 10, C_{31} wax ester; 11, C_{32} *iso*-wax ester; 12, C_{32} wax ester; 13, C_{33} *iso*-wax ester; 14, C_{33} wax ester; 15, C_{34} *iso*-wax ester; 16, C_{34} wax ester; I.S., internal standard.

triacontatriene, the bulk cell material, the fatty acids, and wax esters. This confirms earlier observations of *Chloroflexus* photoautotrophy in these high-sulfide, *Chloroflexus*-dominated microbial mats (12, 39) and shows that carbon isotopic observa-

tions made in culture experiments can be extended to environmental settings.

Chloroflexus biomarkers showed much larger fractionation relative to DIC in the two high-carbonate mats (NMA down-

TABLE 2. Relative abundance of different lipids within mat samples

Category or compound	Peak no. ^a	Relative abundance ^b						
		NMA source ^c	NMA downstream ^d	Tangerine Spring	Octopus Spring	Mushroom Spring	<i>C. aurantiacus</i> ^d	<i>R. castenholzii</i> ^e
<i>Chloroflexus</i> biomarkers								
C _{31:3} alkene	9	++	+	+	Trace	Trace	++	–
<i>n</i> -C ₃₁ wax ester	10	++	+	+	++	++	–	
<i>iso</i> -C ₃₁ wax ester		+	+	+	+	+		
<i>n</i> -C ₃₂ wax ester	12	+++	++	++	+++	+++	+	
<i>iso</i> -C ₃₂ wax ester	11	+	+	+	+	+		
C _{32:1} wax ester							+	
<i>n</i> -C ₃₃ wax ester	14	+++	++	++	+++	+++	+	
<i>iso</i> -C ₃₃ wax ester	13	+	+	+	+	+		
<i>n</i> -C ₃₄ wax ester	16	+++	++	++	+++	+++	++	
<i>iso</i> -C ₃₄ wax ester	15	+	+	+	+	+		
C _{34:1} wax ester							+	
<i>n</i> -C ₃₅ wax ester		+	+	+	++	++	++	
<i>n</i> -C ₃₆ wax ester		+	–	+	+	+	++	
C _{36:1} wax ester							+	
<i>n</i> -C ₃₇ wax ester		+	–	+	+	+	+	+
<i>n</i> -C ₃₈ wax ester								++
<i>n</i> -C ₃₉ wax ester								+
<i>n</i> -C ₄₀ wax ester								++
Cyanobacterial biomarker								
C ₁₇ <i>n</i> -alkane	1	–	++	++	+	+	–	–
Nondiagnostic lipids								
C ₁₅ fatty acid	2	++	++	+	+	++	+	–
C ₁₆ fatty acid	3	++	+++	+++	+	++	+++	–
C ₁₇ fatty acid	4	+	+	+	+	++	+++	–
C ₁₇ alkanol	5	+++	++	++	++	+++	+	–
C _{18:1} fatty acid	6	–	+++	++	–	–	+++	–
C ₁₈ fatty acid	7	+	++	++	+	+	+++	–
C ₁₈ alkanol	8	+++	+	++	+	++	++	–

^a See Fig. 1.^b +, detected; ++, ~50% of base peak; +++, base peak; trace; detected in very low abundance; –, not detected.^c Previously reported by van der Meer et al. (39).^d Knudsen et al. (17), Shiea et al. (35), and van der Meer et al. (40).^e van der Meer et al. (41).

stream and Tangerine Spring) where they live together with cyanobacteria (Fig. 2). This indicates that *Chloroflexus*-like organisms are not purely autotrophic in these cyanobacterial mats. Some isotopically lighter carbon must be obtained from an alternative source. The large $\Delta\delta^{13}\text{C}_{\text{DIC}}$ for the cyanobacterial biomarker C₁₇ *n*-alkane suggests cyanobacteria as a possible source for this light carbon. We considered whether cyanobacterial glucose synthesis and fermentation coupled with cross-feeding could explain the observed isotopic fractionation patterns. The $\Delta\delta^{13}\text{C}_{\text{DIC}}$ of glucose in these two high-carbonate mats is smaller than that observed for the cyanobacterial biomarker, C₁₇ *n*-alkane, consistent with the possibility that cyanobacterial glucose biosynthesis imparts a heavier isotopic signature than lipid biosynthesis (7, 43). However, even in the unlikely event that all of the glucose detected was from cyanobacteria and *Chloroflexus* derived all of its carbon from cyanobacterial sugar fermentation, the $\delta^{13}\text{C}$ value for all *Chloroflexus* biomarkers should be lower than the $\delta^{13}\text{C}$ of glucose, due to isotopic effects of subsequent lipid biosynthesis pathways (1, 14, 20). The fact that the $\delta^{13}\text{C}$ value of the C_{31:3} alkene is higher than (NMA downstream) or similar to (Tangerine Spring) that of glucose and that the $\delta^{13}\text{C}$ value of wax esters (NMA downstream) is similar to that of glucose therefore

suggests that cyanobacterial cross-feeding alone is unlikely to explain the observed results. Apparently, photoautotrophy is also occurring in *Chloroflexus* spp. in these mats. This could reflect either the inputs of separate heterotrophic and autotrophic *Chloroflexus* populations or mixotrophic carbon metabolism in a single *Chloroflexus* population. Since both cyanobacteria and *Chloroflexus*-like organisms contribute organic compounds to the mats, the $\delta^{13}\text{C}$ values for the bulk biomass and glucose must be intermediate to cyanobacterial and autotrophic *Chloroflexus* isotope signatures.


Relative to the Mammoth Terraces cyanobacterial mats, the two mats in alkaline siliceous springs show smaller $\Delta\delta^{13}\text{C}_{\text{DIC}}$ values (Fig. 2). The fractionation pattern of the bulk biomass, glucose, *Chloroflexus* biomarkers, and more general lipids resembles that of the autotrophically grown *C. aurantiacus* culture and the NMA source pool mat, suggesting the possibility of photoautotrophic metabolism by *Chloroflexus* relatives. However, in these mats the $\delta^{13}\text{C}$ values of glucose are sufficiently heavy to support the hypothesis that cyanobacterial sugar biosynthesis imparts a heavier isotopic signature to sugars than to lipids; cross-feeding of sugar fermentation products could then impart the heavier signatures of *Chloroflexus* biomarkers. A complicating factor in Octopus Spring and Mush-

TABLE 3. Stable carbon isotopic compositions of bicarbonate, CO₂ (calculated from bicarbonate), bulk biomass, *Chloroflexus* and cyanobacterial biomarkers, nondiagnostic lipids, and glucose in mat samples expressed in per mille relative to the PeeDee Belemnite standard

Category or compound	Peak no. ^a	$\delta^{13}\text{C}$ values of samples					
		NMA source ^b	NMA downstream ^b	Tangerine Spring	Octopus Spring	Mushroom Spring	<i>C. aurantiacus</i> ^c
Bicarbonate		-2 to -4 ^d	-2 to -4 ^d	-3.3	~6.8 ^e	-6.8	0
CO ₂ ^f		~9	~9	~9	~12	~11	
Bulk biomass		-14.9	-23.5	-24.8	-16.9	-13.2	-13.0
<i>Chloroflexus</i> biomarkers							
C _{31:3} alkene	9	-8.9	-15.1	-20.0	-18.9 ^g	NM ^h	-12.5
C ₃₂ wax ester	12	-17.7	-22.7	-26.0	-18.3	-16.5	NM
C ₃₃ wax ester	14	-17.9	-24.1	-27.0	-18.5	-16.4	-15.0
C ₃₄ wax ester	16	-17.4	-23.4	-26.7	-18.5	-16.5	-14.4
Cyanobacterial biomarker							
C ₁₇ n-alkane	1		-36.3	-36.3	-34.1	-29.6	
General lipids							
C ₁₅ fatty acid	2	-19.4	-23.4	-27.1	-20.2	-17.5	NM
C ₁₆ fatty acid	3	-19.3	-34.5	-35.3	-21.3	-21.3	-13.8
C ₁₇ fatty acid	4	-18.9	-22.1	-27.6	-18.7	-16.8	-14.8
C ₁₇ alkanol	5	-18.0	-22.0	-27.9	-19.4	-15.5	NM
C _{18:1} fatty acid	6		-32.7	-35.5			-12.5
C ₁₈ fatty acid	7	-16.0	-33.9	-35.1	NM	-24.4	-13.6
C ₁₈ alkanol	8	-16.1	-24.3	-29.1	-22.2	-21.5	-12.5
Sugar							
Glucose		-5.1	-21.8	-20.0	-11.0	-11.5	-5.7

^a See Figure 1.^b Previously reported by van der Meer et al. (39), except for the glucose isotope values.^c From van der Meer et al. (40). Isotope values for organic compounds are reported relative to bicarbonate.^d Madigan et al. (19).^e Assuming similar $\delta^{13}\text{C}$ value as for Mushroom Spring.^f Calculated based on temperature-dependent isotope equilibrium equation (21).^g Partial coelution.^h NM, not measured.

room Spring could be the effect of CO₂ limitation on stable carbon isotope fractionation by cyanobacteria in these much more alkaline and lower-DIC settings (8). The Mammoth Terraces mats occur in carbonate-depositing springs that are high in DIC and, at pH 6.4, are poised near the pK_a of H₂CO₃/HCO₃⁻ (i.e., CO₂ is readily available). In contrast, the mats in alkaline silicious springs of the Lower Geyser Basin have midday pHs as high as 9.4, approaching the pK_a of HCO₃⁻/CO₃²⁻ (29). CO₂ limitation in these alkaline silicious hot springs may decrease the degree of isotopic fractionation by cyanobacteria, resulting in higher $\delta^{13}\text{C}$ values for all organic compounds produced by cyanobacteria, including the C₁₇ n-alkane and glucose. However, the very large differentials in $\delta^{13}\text{C}$ values of C₁₇ n-alkane and glucose (18 to 22.4‰) exceed those observed so far in Calvin cycle organisms (1 to 16‰ [43]). This large isotopic difference between the cyanobacterial biomarker and glucose, especially in combination with very small or no isotopic fractionation between CO₂ and glucose, makes it unlikely that all of the glucose is derived from cyanobacteria even when the possible effect of CO₂ limitation on the stable carbon isotope ratios of cyanobacterial products is considered. The high abundance of *Chloroflexus* biomarkers (i.e., wax esters) relative to cyanobacterial biomarkers (i.e., C₁₇ n-alkane) (Fig. 1) and the heavier isotopic signatures of C₁₆ and C₁₈ fatty acids (Table 3), which could be contributed by either type of phototroph (10, 17, 35, 40), indeed suggest that a large fraction of

FIG. 2. Depletions in ¹³C of biomass, key biomarkers, and glucose relative to DIC. \triangleright , biomass; \diamond , hentriacontatriene; \circ , wax esters (average); \times , C₁₇ n-alkane; \square , glucose.

the total biomass might be *Chloroflexus* derived. This might be due to our analysis of thicker mat samples (i.e., >1 cm for Lower Geyser Basin mats versus 1 to 2 mm for Mammoth Terraces mats) and the persistence of *Chloroflexus* carbohydrates in deeper layers of the mats in alkaline siliceous springs (39).

By comparing isotopic compositions of compound classes from replicate cyanobacterial mats from different hot-spring settings, we were able to observe that both autotrophic and heterotrophic carbon metabolisms are employed by *Chloroflexus*-like bacteria. Heavier isotopic signatures of *Chloroflexus*-like bacteria may be due in part to their unique autotrophic biochemistry and in part to the differences in isotopic fractionation in sugar and lipid biosynthetic pathways of cyanobacteria. Further work will be necessary, however, in order to observe the degree to which cyanobacterial sugar biosynthesis affects isotopic compositions and, thus, the relative importance of heterotrophy (via cross-feeding from cyanobacteria) and autotrophy in the carbon metabolism of *Chloroflexus*-like bacteria in these mats. This is especially true of mats in alkaline siliceous springs, where CO₂ limitation effects make resolution of the two types of carbon metabolism more difficult.

ACKNOWLEDGMENTS

We thank M. M. Bateson and B. Lindstrom for help with field work and R. Pancost, B. E. van Dongen, W. I. C. Rijpstra, M. Baas, R. Kloosterhuis, and J. Ossebaar for analytical assistance. We thank The U.S. National Park Service for permission to conduct research in Yellowstone National Park.

This study was supported by U.S. National Aeronautics and Space Administration grants NAGW-2764 and NAG5-3652 and a PIONIER grant awarded to J. S. Sinninghe Damsté by The Netherlands Organization for Scientific Research (NWO). We also thank NWO for supporting travel of M. T. J. van der Meer. We thank Shell International Petroleum Maatschappij BV for financial support for the GC-IRMS facility.

REFERENCES

1. Abraham, W. R., C. Hesse, and O. Pelz. 1998. Ratios of carbon isotopes in microbial lipids as an indicator of substrate usage. *Appl. Environ. Microbiol.* **64**:4202–4209.
2. Anderson, K. L., T. A. Tayne, and D. M. Ward. 1987. Formation and fate of fermentation products in hot-spring cyanobacterial mats. *Appl. Environ. Microbiol.* **53**:2343–2352.
3. Bauld, J., and T. D. Brock. 1973. Ecological studies of *Chloroflexus*, a gliding photosynthetic bacterium. *Arch. Microbiol.* **92**:267–284.
4. Boomer, S. M., D. P. Lodge, B. E. Dutton, and B. I. Pierson. 2002. Molecular characterization of novel red green nonsulfur bacteria from five distinct hot spring communities in Yellowstone National Park. *Appl. Environ. Microbiol.* **68**:346–355.
5. Castenholz, R. W. 1973. The possible photosynthetic use of sulfide by the filamentous phototrophic bacteria of hot springs. *Limnol. Oceanogr.* **18**:863–876.
6. Cline, J. D. 1969. Spectrophotometric determination of hydrogen sulfide in natural waters. *Limnol. Oceanogr.* **14**:454–458.
7. Deines, P. 1980. The isotopic composition of reduced organic carbon, p. 329–406. *In* P. Fritz and J. C. Fontes (ed.), *Handbook of environmental isotope geochemistry*. Elsevier Scientific Publishing Company, Amsterdam, The Netherlands.
8. DesMarais, D. J., J. Bauld, A. C. Palmisano, R. E. Summons, and D. M. Ward. 1992. The biogeochemistry of carbon in modern microbial mats, p. 299–308. *In* J. W. Schopf and C. Klein (ed.), *The proterozoic biosphere: a multidisciplinary study*. Cambridge University Press, Cambridge, United Kingdom.
9. Dobson, G., D. M. Ward, N. R. Robinson, and G. Eglinton. 1988. Biogeochemistry of hot spring environments: free lipids of a cyanobacterial mat. *Chem. Geol.* **68**:155–179.
10. Fork, D. C., N. Murata, and N. Sato. 1979. Effect of growth temperature on the lipid and fatty acid composition, and the dependence on temperature of light-induced redox reactions of cytochrome *f* and of light energy redistribution in the thermophilic blue-green alga *Synechococcus lividus*. *Plant Physiol.* **63**:524–530.
11. Fry, B., W. Brand, F. J. Mensch, K. Tholke, and R. Garritt. 1992. Automated analysis system for coupled $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements. *Anal. Chem.* **64**:288–291.
12. Giovannoni, S. J., N. P. Revsbech, D. M. Ward, and R. W. Castenholz. 1987. Obligately phototrophic *Chloroflexus*: primary production in anaerobic hot spring microbial mats. *Arch. Microbiol.* **147**:80–87.
13. Hanada, S., S. Takaichi, K. Matsuura, and K. Nakamura. 2002. *Roseiflexus castenholzii* gen. nov., sp. nov., a thermophilic, filamentous, photosynthetic bacterium which lacks chlorosomes. *Int. J. Syst. Evol. Microbiol.* **52**:187–193.
14. Hayes, J. M. 2001. Fractionation of carbon and hydrogen isotopes in biosynthetic processes, p. 225–277. *In* J. W. Valley and D. Cole (ed.), *Stable isotope geochemistry. Reviews in mineralogy and geochemistry*, vol. 43. Mineralogical Society of America, Washington, D.C.
15. Herter, S., G. Fuchs, A. Bacher, and W. Eisenreich. 2002. A bicyclic autotrophic CO₂ fixation pathway in *Chloroflexus aurantiacus*. *J. Biol. Chem.* **277**:20277–20283.
16. Holo, H., and R. Sirevåg. 1986. Autotrophic growth and CO₂ fixation of *Chloroflexus aurantiacus*. *Arch. Microbiol.* **145**:173–180.
17. Knudsen, E., E. Jantzen, K. Bryn, J. G. Ormerod, and R. Sirevåg. 1982. Quantitative and structural characteristics of lipids in *Chlorobium* and *Chloroflexus*. *Arch. Microbiol.* **132**:149–154.
18. Konopka, A. 1992. Accumulation and utilization of polysaccharide by hot-spring phototrophs during a light-dark transition. *FEMS Microb. Ecol.* **102**:27–32.
19. Madigan, M. T., R. Takigiku, R. G. Lee, H. Gest, and J. M. Hayes. 1989. Carbon isotope fractionation by thermophilic phototrophic sulfur bacteria: evidence for autotrophic growth in natural populations. *Appl. Environ. Microbiol.* **55**:639–644.
20. Monson, K. D., and J. M. Hayes. 1982. Carbon isotopic fractionation in the biosynthesis of bacterial fatty acids. Ozonolysis of unsaturated fatty acids as a means of determining the intramolecular distribution of carbon isotopes. *Geochim. Cosmochim. Acta* **46**:139–149.
21. Mook, W. G., J. C. Bommerson, and W. H. Staberman. 1974. Carbon isotope fractionation between dissolved bicarbonate and gaseous carbon dioxide. *Earth Planet Sci. Lett.* **22**:169–176.
22. Nold, S. C., and D. M. Ward. 1996. Photosynthate partitioning and fermentation in hot spring microbial mat communities. *Appl. Environ. Microbiol.* **62**:4598–4607.
23. Nübel, U., M. M. Bateson, V. Vandieken, M. Kühl, and D. M. Ward. 2002. Microscopic examination of distribution and phenotypic properties of phylogenetically diverse *Chloroflexaceae*-related bacteria in hot spring microbial mats. *Appl. Environ. Microbiol.* **68**:4593–4603.
24. Oyaizu, H., B. Debrunner-Vossbrinck, L. Mandelco, J. A. Studier, and C. R. Woese. 1987. The green non-sulfur bacteria: a deep branching in the eubacterial line of descent. *Syst. Appl. Microbiol.* **9**:47–53.
25. Pierson, B. K., and R. W. Castenholz. 1974. Studies of pigments and growth in *Chloroflexus aurantiacus*, a phototrophic filamentous bacterium. *Arch. Microbiol.* **100**:283–305.
26. Pierson, B. K., and R. W. Castenholz. 1992. The family Chloroflexaceae, p. 3754–3774. *In* A. Balows, H. G. Trüper, M. Dworkin, W. Harder, and K.-H. Schleifer (ed.), *The prokaryotes*. Springer-Verlag, New York, N.Y.
27. Popp, B. N., E. A. Laws, R. R. Bridgier, J. E. Dore, K. L. Hanson, and S. G. Wakeham. 1998. Effect of phytoplankton cell geometry on carbon isotopic fractionation. *Geochim. Cosmochim. Acta* **62**:69–77.
28. Ramsing, N. B., M. J. Ferris, and D. M. Ward. 2000. Highly ordered vertical structure of *Synechococcus* populations within the one-millimeter-thick photic zone of a hot spring cyanobacterial mat. *Appl. Environ. Microbiol.* **66**:1038–1049.
29. Revsbech, N. P., and D. M. Ward. 1984. Microelectrode studies of interstitial water chemistry and photosynthetic activity in a hot spring microbial mat. *Appl. Environ. Microbiol.* **48**:270–275.
30. Ruff-Roberts, A. L., J. G. Kuennen, and D. M. Ward. 1994. Distribution of cultivated and uncultivated cyanobacteria and *Chloroflexus*-like bacteria in hot spring microbial mats. *Appl. Environ. Microbiol.* **60**:697–704.
31. Sakata, S., J. M. Hayes, A. R. McTaggart, R. A. Evans, K. J. Leckrone, and R. K. Togasaki. 1997. Carbon isotopic fractionation associated with lipid biosynthesis by a cyanobacterium: relevance for interpretation of biomarker records. *Geochim. Cosmochim. Acta* **61**:5379–5389.
32. Sandbeck, K. A., and D. M. Ward. 1981. Fate of immediate methane precursors in low sulfate hot spring algal-bacterial mats. *Appl. Environ. Microbiol.* **41**:775–782.
33. Schouten, S., W. C. M. Klein Breteler, P. Blokker, N. Schogt, W. I. C. Rijpstra, K. Grice, M. Baas, and J. S. Sinninghe Damsté. 1998. Biosynthetic effects on the stable carbon isotopic composition of algal lipids: implications for deciphering the carbon isotopic biomarker record. *Geochim. Cosmochim. Acta* **62**:1397–1406.
34. Shiea, J., S. C. Brassell, and D. M. Ward. 1990. Mid-chain branched mono- and dimethyl alkanes in hot spring cyanobacterial mats: a direct biogenic source for branched alkanes in ancient sediments? *Org. Geochem.* **15**:223–231.
35. Shiea, J., S. C. Brassell, and D. M. Ward. 1991. Comparative analysis of

- extractable lipids in hot spring microbial mats and their component photosynthetic bacteria. *Org. Geochem.* **17**:309–319.
36. **Simon, H., and H. G. Floss.** (1967) Bestimmung der Isotopenverteilung in markierten Verbindungen. Springer, Berlin, Germany.
37. **Summons, R. E., L. L. Jahnke, and B. R. T. Simoneit.** 1996. Lipid biomarkers for bacterial ecosystems: studies of cultured organisms, hydrothermal environments and ancient sediments, p. 174–194. *In* G. R. Block and J. A. Goode (ed.), Evolution of hydrothermal ecosystems on Earth (and Mars?). Ciba Foundation Symposium no. 202. Wiley, Chichester, United Kingdom.
38. **Van der Meer, M. T. J., S. Schouten, D. M. Ward, J. A. J. Geenevasen, and J. S. Sinninghe Damsté.** 1999. All-cis hentriaconta-9,15,22-triene in microbial mats formed by the phototrophic prokaryote *Chloroflexus*. *Org. Geochem.* **30**:1585–1587.
39. **Van der Meer, M. T. J., S. Schouten, J. W. de Leeuw, and D. M. Ward.** 2000. Autotrophy of green non-sulphur bacteria in hot spring microbial mats: biological explanations for isotopically heavy organic carbon in the geological record. *Env. Microbiol.* **2**:428–435.
40. **Van der Meer, M. T. J., S. Schouten, B. E. van Dongen, W. I. C. Rijpstra, G. Fuchs, J. S. Sinninghe Damsté, J. W. de Leeuw, and D. M. Ward.** 2001. Biosynthetic controls on the ^{13}C -contents of organic components in the photoautotrophic bacterium *Chloroflexus aurantiacus*. *J. Biol. Chem.* **276**:10971–10976.
41. **Van der Meer, M. T. J., S. Schouten, S. Hanada, E. C. Hopmans, S. Sinninghe Damsté, and D. M. Ward.** 2002. Alkane-1,2-diol-based glycosides and fatty glycosides and wax esters in *Roseiflexus castenholzii* and hot spring microbial mats. *Arch. Microbiol.* **178**:229–237.
42. **Van Dongen, B. E., S. Schouten, and J. S. Sinninghe Damsté.** 2001. Gas chromatography/combustion/isotope-ratio-monitoring mass spectrometric analysis of methylboronic derivatives of monosaccharides: a new method for determining natural ^{13}C abundances of carbohydrates. *Rapid Commun. Mass Spectrom.* **15**:496–500.
43. **Van Dongen, B. E., S. Schouten, and J. S. Sinninghe Damsté.** 2002. Carbon isotopic variability in algal and terrestrial carbohydrates. *Mar. Ecol. Prog. Ser.* **232**:83–92.
44. **Ward, D. M., T. A. Tayne, K. L. Anderson, and M. M. Bateson.** 1987. Community structure, and interactions among community members in hot spring cyanobacterial mats. *Symp. Soc. Gen. Microbiol.* **41**:179–210.
45. **Ward, D. M., R. Weller, J. Shiea, R. W. Castenholz, and Y. Cohen.** 1989. Hot spring microbial mats: anoxygenic and oxygenic mats of possible evolutionary significance, p. 3–15. *In* Y. Cohen and E. Rosenberg (ed.), Microbial mats: physiological ecology of benthic microbial communities. American Society for Microbiology, Washington, D.C.
46. **Ward, D. M., C. M. Santegoeds, S. C. Nold, N. B. Ramsing, M. J. Ferris, and M. M. Bateson.** 1997. Biodiversity within hot spring microbial mat communities: molecular monitoring of enrichment cultures. *Antonie Leeuwenhoek* **71**:143–150.
47. **Ward, D. M., M. J. Ferris, S. C. Nold, and M. M. Bateson.** 1998. A natural view of microbial biodiversity within hot spring cyanobacterial mat communities. *Microbiol. Mol. Biol. Rev.* **62**:1353–1370.